arXiv:1606.09569v1 [quant-ph] 29 Jun 2016

Green's function of a general PT-symmetric non-Hermitian non-central potential

Brijesh Kumar Mourya and Bhabani Prasad Mandal

Abstract We study the path integral solution of a system of particle moving in certain class of PT symmetric non-Hermitian and non-central potential. The Hamiltonian of the system is converted to a separable Hamiltonian of Liouville type in parabolic coordinates and is further mapped into a Hamiltonian corresponding to two 2- dimensional simple harmonic oscillators (SHOs). Thus the explicit Green's functions for a general non-central PT symmetric non hermitian potential are calculated in terms of that of 2d SHOs. The entire spectrum for this three dimensional system is shown to be always real leading to the fact that the system remains in unbroken PT phase all the time

.

1 Introduction

Feynman's path integral (PI) approach to quantum mechanical systems is an elegant formalism and powerful in semi-classical calculations [1]. PI formalism which is generally tied to the Lagrangian formalism of mechanics is an extremely powerful technique in quantum mechanics. In a class of problems it provides the Green's function with tremendous ease and also provides valuable insight into the relation between classical and quantum mechanics. Green's functions which in mathematics are to solve non-homogeneous boundary value problems are the backbone of any calculations of physical quantities in quantum field theory [2]. Thus it is extremely important for any physical theory that the Green's functions are well defined. The

Brijesh Kumar Mourya
Department of Physics, Banaras Hindu University, Varanasi-221005,India.e-mail: brijeshkumarbhu@gmail.com

Bhabani Prasad Mandal
Department of Physics, Banaras Hindu University, Varanasi-221005,India. e-mail: bhabani.mandal@gmail.com

purpose of the present work is to discuss the PI formulation of a general non-central, combined parity (P) and time reversal (T) symmetric non-Hermitian system in 3d by calculating the explicit Green's functions for such a system. Consistent quantum theory with real energy eigenvalues, unitary time evolution and probabilistic interpretation for PT symmetric non-Hermitian theories in a different Hilbert space equipped with positive definite inner product has been the subject of intrinsic research in frontier physics over the last one and half decades [3]. Such non-Hermitian PT symmetric systems generally exhibit PT phase transition or more specifically a PT breaking transition [4] which has been realised experimentally [5]. In-spite of huge success and wide applicability [6, 7, 8, 9, 10, 11] of this field the study of non-Hermitian quantum mechanics is mostly restricted to one dimensional or central potentials in higher dimension potential problems. In this article we consider a general and very important non-central PT symmetric non-Hermitian system in the PI formulation. First we show that how the Hamiltonian corresponding to this potential is reduced to a separable Hamiltonian of Liouville type [12] in a different coordinate system. This further enables us to map the system into two non-interacting 2d harmonic oscillators with the appropriate choice of coordinates. We then calculate the Green's functions of the system in terms of the Green's functions of 2d harmonic oscillators. Further we write the Hamiltonian in terms of appropriate creation and annihilation operators to calculate the energy eigenvalues of this non-central non-Hermitian system. We find that the energy eigenvalues are always real as long as the parameters in the potential are real. This indicates that system is always in unbroken PT phase.

Now we present the plan of the paper. In Sec. 2 we calculate the Green's function for the system in terms of SHO Green's functions. The reality of the spectrum is shown in Sec. 3. Sec. 4 is kept for concluding remarks.

2 Green's functions for the PT-symmetric non-central potential

We consider a system described by a general non-central non-Hermitian potential in 3-dimension in spherical polar coordinates as

$$H = \frac{P_r^2}{2m} + \frac{P_\theta^2}{2mr^2} + \frac{P_\phi^2}{2mr^2 sin^2\theta} + V(r,\theta) \tag{1}$$

where the non-Hermitian potential is

$$V(r,\theta) = -\frac{\alpha}{r} + \frac{B\hbar^2}{2mr^2\sin^2\theta} + \frac{iC\hbar^2\cos\theta}{2mr^2\sin^2\theta}. \tag{2}$$

α, B and C are real constants. It is straight forward to check that this non-Hermitian system is PT symmetric, where in 3-d in spherical polar coordinates the parity transformation is defined as, $r \to r; \quad \theta \to \pi - \theta, \quad \phi \to \phi + 2\pi$.This particular potential is very important as the Coulomb and the ring- shaped potentials are particular cases

of this potential. For $C = 0$ this potential becomes Hartman's ring shaped potential which was originally proposed to model Benzene molecule [14]. To proceed with this Hamiltonian we first consider the most general Hamiltonian of Liouville type written as

$$H = \frac{1}{V_1(q_1)+V_2(q_2)}\left\{\frac{1}{2mW_1(q_1)}p_1^2+\frac{1}{2mW_2(q_2)}p_2^2+U_1(q_1)+U_2(q_2)\right\} \quad (3)$$

This is reduced to a simpler form by using some canonical transformation and re-defining U 's and V's as [12]

$$H = \frac{1}{V_1(q_1)+V_2(q_2)}\left\{\frac{1}{2m}p_1^2+\frac{1}{2m}p_2^2+U_1(q_1)+U_2(q_2)\right\} \quad (4)$$

The time independent Schrodinger equation corresponding to this Hamiltonian is then written as

$$\hat{H}_T\psi = 0$$

with a total Hamiltonian is defined as

$$\hat{H}_T = \frac{1}{2m}p_1^2+\frac{1}{2m}p_2^2+U_1(q_1)+U_2(q_2)-E(V_1(q_1)+V_2(q_2)) \quad (5)$$

Now, to obtain the path integral for this Hamiltonian $\hat{H}_T$, let us consider the evaluation of this operator for a arbitrary parameter τ :

$$\begin{aligned}\left\langle q_{1b},q_{2b}\left|\exp\left\{-i\frac{\hat{H}_T\tau}{\hbar}\right\}\right|q_{1a},q_{2a}\right\rangle &= \left\langle q_{1b}\left|\exp\left\{-(\frac{i}{\hbar})\left[\frac{1}{2m}\hat{p}_1^2+U_1(q_1)-EV_1(q_1)\right]\tau\right\}\right|q_{1a}\right\rangle \\ &\times\left\langle q_{2b}\left|\exp\left\{-(\frac{i}{\hbar})\left[\frac{1}{2m}\hat{p}_2^2+U_2(q_2)-EV_2(q_2)\right]\tau\right\}\right|q_{2a}\right\rangle\end{aligned} \quad (6)$$

The RHS of Eq. 6 is written in terms of path integral [12].

$$\begin{aligned}&\left\langle q_{1b},q_{2b}\left|\exp\left\{-i\frac{\hat{H}_T\tau}{\hbar}\right\}\right|q_{1a},q_{2a}\right\rangle \\ &= \int\mathscr{D}q_1\mathscr{D}p_1\exp\left\{(\frac{i}{\hbar}\int_0^\tau\left[p_1\dot{q}_1-\left(\frac{\hat{p}_1^2}{2m}+U_1(q_1)-EV_1(q_1)\right)\right]d\tau\right\} \\ &\times\int\mathscr{D}q_2\mathscr{D}p_2\exp\left\{(\frac{i}{\hbar}\int_0^\tau\left[p_2\dot{q}_2-\left(\frac{\hat{p}_2^2}{2m}+U_2(q_2)-EV_2(q_2)\right)\right]d\tau\right\}\end{aligned} \quad (7)$$

The parameter τ is arbitrary and one can obtain physically meaningful quantity out of Eq. 6 by integrating over τ from 0 to ∞ .

$$\left\langle q_{1b},q_{2b}\left|\frac{\hbar}{\hat{H}_T}\right|q_{1a},q_{2a}\right\rangle_{\text{Semi-classical}}$$

$$= \int_0^\infty d\tau \left\langle q_{1b} \left| \exp\left\{ -(\frac{i}{\hbar}) \left[\frac{1}{2m}\hat{p}_1^2 + U_1(q_1) - EV_1(q_1) \right] \tau \right\} \right| q_{1a} \right\rangle$$
$$\times \left\langle q_{2b} \left| \exp\left\{ -(\frac{i}{\hbar}) \left[\frac{1}{2m}\hat{p}_2^2 + U_2(q_2) - EV_2(q_2) \right] \tau \right\} \right| q_{2a} \right\rangle \tag{8}$$

The meaning of RHS of Eq. 8 has been explained in [12]. And the LHS of it is written as,

$$\left\langle q_{1b}, q_{2b} \left| \frac{\hbar}{\hat{H}_T} \right| q_{1a}, q_{2a} \right\rangle = \left\langle q_{1b}, q_{2b} \left| \frac{\hbar}{\hat{H} - E} \right| q_{1a}, q_{2a} \right\rangle \frac{1}{V_1(q_{1a}) + V_2(q_{2a})} \tag{9}$$

with the completeness relation

$$\int |q_1\, q_2\rangle \frac{dV}{V_1(q_1) + V_2(q_2)} \langle q_1\, q_2| = 1 \tag{10}$$

Now we show that the Hamiltonian of a particle in a non-central potential is reduced to a separable Hamiltonian of the above kind. Let us start with the non-central system written in Eq. 2

$$H = \frac{P_r^2}{2m} + \frac{P_\theta^2}{2mr^2} + \frac{P_\phi^2}{2mr^2 sin^2\theta} - \frac{\alpha}{r} + \frac{\beta}{r^2 \sin^2\theta} + \frac{i\gamma\cos\theta}{r^2 \sin^2\theta} \tag{11}$$

where $\beta = \frac{B\hbar^2}{2m}$; $\gamma = \frac{C\hbar^2}{2m}$. This system will be reduced to separable system of Liouville type in parabolic coordinate system. To express the Hamiltonian in parabolic coordinates (ξ, η, ϕ), it is useful to first express this potential $V(r,\theta)$ in cylindrical coordinates (ρ, ϕ, z). In cylindrical coordinate the potential looks like,

$$V(\rho, z) = -\frac{\alpha}{\sqrt{\rho^2 + z^2}} + \frac{\beta}{\rho^2} + \frac{i\gamma z}{\rho^2\sqrt{\rho^2 + z^2}}; \quad \rho^2 = x^2 + y^2 \tag{12}$$

The parabolic coordinates are expressed in terms of cylindrical coordinates as

$$\xi = \frac{1}{2}\left(\sqrt{\rho^2 + z^2} - z\right);$$
$$\eta = \frac{1}{2}\left(\sqrt{\rho^2 + z^2} + z\right); \quad \phi = \phi\ . \tag{13}$$

Now the potential in Eq. 12 in terms of these parabolic coordinates, is

$$V(\xi, \eta) = -\frac{\alpha}{\xi + \eta} + \frac{\beta}{4\xi\eta} + \frac{i\gamma(\eta - \xi)}{4\eta\xi(\eta + \xi)} \tag{14}$$

and the Hamiltonian in parabolic coordinate is written as

$$H(\xi,\eta,\phi)=\frac{1}{2m(\xi+\eta)}\left[\xi\hat{p}_\xi^2+\eta\hat{p}_\eta^2\right]+\frac{1}{8m\eta\xi}\hat{p}_\phi^2+V(\xi,\eta) \tag{15}$$

We define variables u,v

$$\begin{aligned}\xi &= \frac{1}{4}u^2; \quad 0\le u<\infty\\ \eta &= \frac{1}{4}v^2; \quad 0\le v<\infty\end{aligned} \tag{16}$$

and perform a canonical transformation

$$\sqrt{\xi}\hat{p}_\xi=\hat{p}_u\ ;\ \sqrt{\eta}\hat{p}_\eta=\hat{p}_v \tag{17}$$

to simplify the kinetic term in H in Eq. 15 as

$$H(u,v,\phi)=\frac{4}{u+v}\left\{\frac{1}{2m}\left[\hat{p}_u^2+\hat{p}_v^2+(\frac{1}{u^2}+\frac{1}{v^2})\hat{p}_\phi^2\right]-\alpha+\frac{\beta+i\gamma}{u^2}+\frac{\beta-i\gamma}{v^2}\right\} \tag{18}$$

This is further written compactly as

$$H(u,v,\phi)=\frac{4}{u^2+v^2}\left\{\frac{1}{2m}\left[\hat{p}_u^2+\hat{p}_v^2+\frac{1}{u^2}\hat{p}_{\phi_1}^2+\frac{1}{v^2}\hat{p}_{\phi_2}^2\right]-\alpha\right\} \tag{19}$$

where

$$\hat{p}_{\phi_1}^2=\hat{p}_\phi^2+2m(\beta+i\gamma);\quad \hat{p}_{\phi_2}^2=\hat{p}_\phi^2+2m(\beta-i\gamma) \tag{20}$$

Note $\hat{p}_{\phi_1}$ and $\hat{p}_{\phi_2}$ are not Hermitian but complex conjugate to each other. This Hamiltonian is still not a separable one of Liouville type. We further consider the total Hamiltonian $H_T(=H-E)$ with $E=-2m\omega^2$ for the bound state case ($E<0$),

$$\hat{H}_T=\frac{1}{2m}\left[\hat{p}_u^2+\hat{p}_v^2+\frac{1}{u^2}\hat{p}_{\phi_1}^2+\frac{1}{v^2}\hat{p}_{\phi_2}^2\right]-\alpha+\frac{1}{2}m\omega^2(u^2+v^2) \tag{21}$$

Now we introduce the components of 2-dimensional vectors **u** and **v** as

$$\begin{aligned}\mathbf{u} &= (u_1,u_2)=(u\cos\phi_1,u\sin\phi_1)\\ \mathbf{v} &= (v_1,v_2)=(v\cos\phi_2,v\sin\phi_2)\end{aligned} \tag{22}$$

to have

$$\mathbf{p_u}^2=\hat{p}_u^2+\frac{1}{u^2}\hat{p}_{\phi_1}^2;\qquad \mathbf{p_v}^2=\hat{p}_v^2+\frac{1}{v^2}\hat{p}_{\phi_2}^2; \tag{23}$$

Putting all these in Eq. 21 we obtain

$$\hat{H}_T=\frac{1}{2m}\mathbf{p}_u^2+\frac{m\omega^2}{2}\mathbf{u}^2+\frac{1}{2m}\mathbf{p}_v^2+\frac{m\omega^2}{2}\mathbf{v}^2-\alpha \tag{24}$$

This is the Hamiltonian which is separable of Liouville type. Thus, the Hamiltonian for the non-central potential has been reduced to that of two 2-dimensional oscillators (apart from some constant shift in ground state energy).

Now by using Eq. 6 for this separable Hamiltonian, we find the path integral for the non-Hermitian non-central potential exactly as

$$\begin{aligned}\left\langle \mathbf{u}_b,\mathbf{v}_b\left|\exp\left[-i\frac{\hat{H}_T\tau}{\hbar}\right]\right|\mathbf{u}_a,\mathbf{v}_a\right\rangle &= e^{\frac{i\alpha\tau}{\hbar}}\left\langle \mathbf{u}_b\left|\exp\left[-(\frac{i}{\hbar})\left(\frac{\mathbf{p}_u^2}{2m}+\frac{m\omega^2}{2}\mathbf{u}^2\right)\tau\right]\right|\mathbf{u}_a\right\rangle \\ &\times\left\langle \mathbf{v}_b\left|\exp\left[-(\frac{i}{\hbar})\left(\frac{\mathbf{p}_v^2}{2m}+\frac{m\omega^2}{2}\mathbf{v}^2\right)\tau\right]\right|\mathbf{u}_a\right\rangle \\ = e^{(\frac{i\alpha\tau}{\hbar})}\left(\frac{im\omega}{2\pi i\hbar\sin\omega\tau}\right)^{\frac{4}{2}}&\exp\left\{\frac{im\omega}{2\hbar\sin\omega\tau}\left[(\mathbf{u}_b^2+\mathbf{v}_b^2+\mathbf{u}_a^2+\mathbf{v}_a^2)\cos\omega\tau-2\mathbf{u}_b\cdot\mathbf{u}_a-2\mathbf{v}_b\cdot\mathbf{v}_a\right]\right\}\end{aligned} \tag{25}$$

where the exact result for one dimensional simple harmonic oscillator has been used [1].

$$\left\langle q_b\left|\exp\left[-\frac{i}{\hbar}\left(\frac{\hat{p}^2}{2m}+\frac{m\omega^2q^2}{2}\right)\tau\right]\right|q_a\right\rangle=\left(\frac{m\omega}{2\pi i\hbar\sin\omega\tau}\right)^{\frac{1}{2}}\exp\left\{\frac{im\omega}{2\hbar\sin\omega\tau}\left[(q_a^2+q_b^2)\cos\omega\tau-2q_aq_b\right]\right\} \tag{26}$$

The Eq. 25 contains the arbitrary parameter τ and has to be eliminated to obtain physically meaningful quantity. This can be done by integrating over τ from 0 to ∞ in both side of the Eq. 25. When we integrate over τ in the LHS of the Eq. 25, it is nothing but the Green's functions of the operator $\frac{1}{\hat{H}-E}$ as discussed at beginning of this section. And the integration in the RHS can be done in a straightforward manner [12]. Thus we obtain the explicit Green's functions for the system of non central non-Hermitian potential.

3 Reality of the Spectrum

Since this system with non-Hermitian, non-central potential is equivalent to two 2d SHOs, conjugate to each other we define the creation and annihilation operators for this theory as

$$\begin{aligned}a_k&=\frac{1}{\sqrt{2}}\left[\sqrt{\frac{m\omega}{\hbar}}u_k+\frac{i}{\sqrt{m\omega\hbar}}\hat{p}_{u_k}\right]\\ \tilde{a}_k&=\frac{1}{\sqrt{2}}\left[\sqrt{\frac{m\omega}{\hbar}}v_k+\frac{i}{\sqrt{m\omega\hbar}}\hat{p}_{v_k}\right]\end{aligned} \tag{27}$$

where $k=1,2$ to obtain the energy levels for this system of non-central potential in a simple algebraic way. The total Hamiltonian in Eq. 24 is written in terms of these

creation and annihilation operators as follows:

$$\hat{H}_T = \hbar\omega\left[\sum_{k=1}^{2}\left(a_k^\dagger a_k + \tilde{a}_k^\dagger \tilde{a}_k\right) + 2\right] - \alpha \tag{28}$$

and the conjugate momentum variables are written as

$$\begin{aligned}\hat{p}_{\phi_1} &= i\hbar\left[a_1^\dagger a_2 - a_2^\dagger a_1\right] \\ \hat{p}_{\phi_2} &= i\hbar\left[\tilde{a}_1^\dagger \tilde{a}_2 - \tilde{a}_2^\dagger \tilde{a}_1\right]\end{aligned} \tag{29}$$

We further perform an unitary transformation of the following type,

$$\begin{aligned}a_1 &= \frac{1}{\sqrt{2}}\left(b_1 - ib_2\right) \\ a_2 &= \frac{1}{\sqrt{2}}\left(-ib_1 + b_2\right)\end{aligned} \tag{30}$$

and similar transformations for $\tilde{a}_1, \tilde{a}_2$ also in Eqs. 28 and 29 to get,

$$\hat{H}_T = \hbar\omega\left[\sum_{k=1}^{2}\left(b_k^\dagger b_k + \tilde{b}_k^\dagger \tilde{b}_k\right) + 2\right] - \alpha \tag{31}$$

and

$$\begin{aligned}\hat{p}_{\phi_1} &= \hbar\left[b_1^\dagger b_1 - b_2^\dagger b_2\right] \\ \hat{p}_{\phi_2} &= \hbar\left[\tilde{b}_1^\dagger \tilde{b}_1 - \tilde{b}_2^\dagger \tilde{b}_2\right]\end{aligned} \tag{32}$$

The number operators, $n_k = b_k^\dagger b_k$, $\tilde{n}_k = \tilde{b}_k^\dagger \tilde{b}_k$ are defined for this system. In terms of number operators the total Hamiltonian in Eq. 31 is now written as

$$\begin{aligned}\hat{H}_T &= \hbar\omega\left[n_1 + n_2 + \tilde{n}_1 + \tilde{n}_2 + 2\right] - \alpha \\ &= 2\hbar\omega\left[n_2 + \tilde{n}_2 + 1 + \frac{\hat{p}_{\phi_1} + \hat{p}_{\phi_2}}{\hbar}\right] - \alpha\end{aligned} \tag{33}$$

Now considering $\left[\hat{p}_{\phi_1} + \hat{p}_{\phi_2}\right]\phi_{phy} \equiv \lambda\,\phi_{phy}$, the physical state condition is [12],

$$\left[2(n_2 + \tilde{n}_2 + 1)\hbar\omega + \omega\lambda - \alpha\right]\phi_{phy} = 0 \tag{34}$$

Hence the energy level can be written in terms of λ as,

$$E_{n_2,\tilde{n}_2,\lambda} \equiv -2m\omega^2 = -\frac{2m\alpha^2}{\left[2(n_2 + \tilde{n}_2 + 1)\hbar + \lambda\right]^2}. \tag{35}$$

λ can be calculated easily using $\left[\hat{p}_{\phi_1} + \hat{p}_{\phi_2}\right]\phi_{phy} \equiv \lambda\,\phi_{phy}$ and Eq. 20 as

$$\lambda = \hbar\left[\left(\nu^2+B+iC\right)^{\frac{1}{2}}+\left(\nu^2+B-iC\right)^{\frac{1}{2}}\right]$$
$$= \left[\left(\nu^2\hbar^2+2m(\beta+i\gamma)\right)^{\frac{1}{2}}+\left(\nu^2\hbar^2+2m(\beta-i\gamma)\right)^{\frac{1}{2}}\right] \quad (36)$$

where ν is non-negative integer and λ is real as $\lambda = \lambda^*$. Therefore, the complete real bound state spectrum for the problem is

$$E_{n_2,\tilde{n}_2,\nu} = \frac{-m\alpha^2}{2\hbar^2\left[n_2+\tilde{n}_2+1+\frac{\sqrt{\nu^2+B+iC}+\sqrt{\nu^2+B-iC}}{2}\right]^2}$$
$$= \frac{-m\alpha^2}{2\left[(n_2+\tilde{n}_2+1)\hbar+\frac{\sqrt{\nu^2\hbar^2+2m(\beta+i\gamma)}+\sqrt{\nu^2\hbar^2+2m(\beta-i\gamma)}}{2}\right]^2} \quad (37)$$

The corresponding result for the real potential agrees with that of in Refs. [15, 16] where energy spectrum has been calculated by solving Schroedinger equation using complicated KS transformation [17, 18].

4 Conclusion

The Hamiltonian corresponding to the PT symmetric non-Hermitian non-central potential in Eq. 2 has been mapped into a Hamiltonian of two 2d harmonic oscillators by choosing appropriate coordinate system and using a suitable canonical transformation. Next we have calculated the Green's functions for the system using path integral method for this separable Hamiltonian of Liouville type. The exact spectrum are calculated by writing this Hamiltonian in terms of creation and annihilation operators of 2d SHO. The entire spectrum is real for any real values of the parameters α, β and γ indicating that system is always in unbroken phase.

Acknowledgments: BPM acknowledge the financial support from the Department of Science and Technology (DST), Govt. of India under SERC project sanction grant No. SR/S2/HEP-0009/2012.

References

1. R.P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integral* (McGraw Hill, New York, 1965)
2. See for example, A Das, *Field Theory: A Path Integral Approach* (World Scientific)
3. C. M. Bender and S. Boettcher *Phys.Rev.Lett.* **80**, 5243 (1998); A. Mostafazadeh, *Int. J. Geom. Meth. Mod. Phys.* **7**, 1191(2010) and references therein; C. M. Bender, *Rep.Prog.*

Phys. **70**, 947 (2007) and references therein.
4. G. Levai, J. Phys. A 41 (2008) 244015; C. M. Bender, G. V. Dunne, P. N. Meisinger, M. Simsek Phys. Lett.A 281 (2001)311-316. C. M. Bender, David J. Weir J. Phys. A 45 (2012)425303; B. P. Mandal, B. K. Mourya, and R. K. Yadav (BHU),Phys. Lett. A 377, 1043 (2013); B. P. Mandal, B. K. Mourya, K, Ali and A. Ghatak, arXiv. 1509.07500 (Press, Ann. of Phys. (2015)); M. Znojil J. Phys.A 36(2003) 7825; C. M. Bender, G. V. Dunne, P. N. Meisinger, M. Simsek Phys. Lett. A 281 (2001)311-316.
5. C. E. Ruter, K. G. Makris, R. El-Ganainy, D. N. Christodulides, M. Segev, and D. Kip, Nature (London) Phys. 6, 192 (2010); A. Guo, G. J. Salamo, Phys. Rev. Lett. 103, 093902 (2009); C. M. Bender, S. Boettcher and P. N. Meisinger, J. Math. Phys. 40 2201 (1999); C. T. West, T. Kottos, and T. Prosen,Phys. Rev. Lett. 104, 054102(2010); A. Nanayakkara Phys. Lett. A 304, 67 (2002)
6. Z. H. Musslimani, K. G. Makris, R. El-Ganainy, and D. N. Christodou lides, Phys. Rev. Lett. 100, 030402 (2008);M. V. Berry, Czech. J. Phys. 54, 1039 (2004); W. D. Heiss,Phys. Rep. 242, 443 (1994).
7. C. M. Bender, D. C. Brody, H. F. Jones, *Phys.Rev. D* **70**, 025001 (2004). C. M. Bender, D. C. Brody, J. Caldeira, B. K. Meister, *arXiv* **1011.1871** (2010). S. Longhi *Phys. Rev. B* **80**, 165125 (2009). B. F. Samsonov *J. Phys. A* **43**, 402006 (2010); *Math. J. Phys. A*: *Math. Gen.* **38**, L571 (2005).
8. B. Basu-Mallick and B.P. Mandal, Phys. Lett. A 284, 231 (2001); B. Basu-Mallick, T. Bhattacharyya and B. P. Mandal Mod. Phys. Lett. A 20, 543 (2004);
9. A. Ghatak, R. D. Ray Mandal, B. P. Mandal,Ann. of Phys.336, 540 (2013); A. Ghatak, J. A. Nathan, B. P. Mandal, and Z. Ahmed,J. Phys. A: Math. Theor.45, 465305 (2012); A. Mostafazadeh, Phys. Rev. A 87, 012103 (2013); L. Deak, T. Fulop Ann. of Phys. 327, 1050 (2012); J. N. Joglekar, C. Thompson, D. D. Scott and G Vemuri, Eur. Phys. J. Appl. Phys. 63, 30001 (2013) ; M. Hasan, A. Ghatak, B. P. Mandal,Ann. of Phys. 344, 17 (2014); A. Ghatak and B. P. Mandal J. Phys. A: Math. Theor. 45, 355301 (2012); S. Longhi, *J. Phys. A: Math. Theor.* **44**, 485302 (2011).
10. B. P. Mandal and S S. Mahajan , Comm. Theor. Phys. 64, 425(2015) ; S Dey and A Fring Phys. Rev A 88, 022116 (2013); C. M. Bender, D. W. Hook, P. N. Meisinger, Q. Wang, *Phys.Rev.Lett.* **104**, 061601 (2010); C. M. Bender, D. W. Hook, K. S. Kooner *J.Phys.A* **43**, 165201, (2010).
11. A. Khare and B. P. Mandal, *Phys.Lett.* **A272**, 53 (2000); B. P. Mandal, S. Gupta, *Mod.Phys.Lett. A* **25**, 1723 (2010); B. P. Mandal, *Mod. Phys. Lett. A* **20**, 655(2005); B. P. Mandal and A. Ghatak, J. Phys. A: Math. Theor. 45, 444022 (2012); M Znojil, Ann. of Phys. 361, 226 (2015).
12. K. Fujikawa, Nucl. Phys. **B 484**, 495, (1997);
13. B P Mandal, Int. J of Mod. Phys A 15, 1225 (2000).
14. H. Hartmann, Theor. Chim. Acta **24** 201 (1972).
15. M. Kibler and C. Campigotto, Int. J. of Quantum Chemistry, **45**, 209(1993).
16. L. Chetouani, L. Guechi and T. F. Hammann, J. Math. Phys. **33** 3410 (1992).
17. L. H. Duru and H. Kleinert, Phys. Lett. **B84**, 185(1979).
18. P. Kustaanheimo and E. Stiefel, J. Reine Angew Math. **218** 204(1965).